

2018 AHP Quality Measures

<u>Measure</u>	<u>Measure Description</u>	<u>90th %ile</u>
Adult Preventive		
Adult BMI	The percentage of patients 18-74 years of age who had an outpatient visit and who had their body mass index (BMI) documented during the measurement year or the year prior the measurement year.	91.73%
Breast Cancer Screening	The percentage of women 50–74 years of age who had a mammogram to screen for breast cancer.	79.84%
Chlamydia Screening	Percentage of women ages 16-24 years who were identified as sexually active who had at least one test for Chlamydia during the measurement year.	79.99%
Colorectal Cancer Screening	The percentage of adults 50–75 years of age who had appropriate screening for colorectal cancer.	72.15%
Fall Risk Mgmt	Percentage of patients 65 years of age or older who had a fall or had problems with balance or walking in the past 12 months, who were seen by a practitioner in the past 12 months and who received fall risk intervention from their current practitioner.	74.00%
Influenza Vaccination (65+)	Percentage of patients 65 years of age and older who received an influenza vaccination during the measurement period.	77.00%
Adult Acute		
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.	39.47%
Use of Imaging Studies for Low Back Pain	The percentage of patients with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis.	82.37%
Medication Reconciliation Post-Discharge	Percentage of discharges of patients 18 years of age or older for whom medications were reconciled on or within 30 days of discharge.	TBD
Osteoporosis Mgmt	Percentage of female members 67 and older who suffered a fracture during the measurement year and who subsequently had either a bone mineral density test or were prescribed a drug to treat or prevent osteoporosis in the six months after the fracture.	71.00%

2018 AHP Quality Measures

Measure	Measure Description	90th %ile
<u>Disease Management</u>		
Medical Attention for Nephropathy	The percentage of patients 18–75 years of age with diabetes (type 1 and type 2) with a nephropathy screening test or evidence of nephropathy, defined as evidence of treatment for nephropathy, evidence of a visit to a nephrologist, a micro albumin test, an ACEI, ARB, or a positive macro albumin test in the measurement year.	92.64%
Diabetes BP Control	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) whose most recent blood pressure in the measurement year was less than 140/90 mmHG.	76.46%
Diabetes Eye Exam	The percentage of patients 18–75 years of age with diabetes (type 1 and type 2) who had a retinal exam or dilated eye exam by an eye care specialist.	67.99%
Diabetes HbA1c <8	The percentage of patients 18–75 years of age with diabetes (type 1 and type 2) whose most recent HbA1c in the measurement year was <8.0%.	65.21%
Diabetes HbA1c<=9	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) whose most recent HbA1c in the measurement year was less than or equal to 9.0%.	77.62%
HTN BP Control	<p>The percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose most recent BP was adequately controlled based on the following criteria:</p> <ul style="list-style-type: none"> - Patients 18-59 years of age whose BP was <140/90 - Patients 60-85 years of age with a diagnosis of diabetes whose BP was <140/90 - Patients 60-85 years of age without a diagnosis of diabetes whose BP was <150/90 	75.18%

2018 AHP Quality Measures

<u>Measure</u>	<u>Measure Description</u>	<u>90th %ile</u>
<u>Pediatric Preventive</u>		
Adolescent Immunizations (Combo 1)	Percentage of adolescents 13 years of age who had 1 meningococcal vaccine and 1 Tdap by their 13th birthday.	87.88%
Adolescent Well Care Visits	Percentage of members 12-21 years of age who had 1 or more well care visit with a PCP or OBGYN during the measurement year.	70.41%
BMI Percentile (Adol)	The percentage of children 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of BMI percentile documentation during the measurement year.	TBD
Childhood Immunizations (Combo 3)	Percentage of children 2 years of age who had 4 DTaP, 3 IPV, 3 Hib, 3 Hep B, 1 MMR, 1 VZV, and 4 pneumococcal conjugate vaccines by their 2nd birthday.	87.04%
Lead Screening	Percentage of children 2 years of age who had one or more capillary or venous blood tests for lead poisoning by their second birthday.	86.37%
Well Care Visits - 3-6 Years	Percentage of members 3-6 years of age who had 1 or more WCV with a PCP during the measurement year.	87.28%
Well Care Visits -First 15 months (6 visits)	Percentage of members who turned 15 months old during the measurement year and who had at least 6 well care visits by their 15 month birthdate.	88.33%
<u>Pediatric Acute</u>		
Appropriate Testing for Children with Pharyngitis	Percentage of children 2-18 years of age who were diagnosed with pharyngitis, dispensed an antibiotic and received group A strep test for the episode.	93.11%
Appropriate Treatment - Children with a URI	Percentage of children 3 months to 18 years of age who were given a diagnosis of URI and were NOT dispensed an antibiotic prescription.	94.80%